




THE THERAPIST
JOURNAL OF THERAPIES & REHABILITATION SCIENCES
<https://thetherapist.com.pk/index.php/tt>
ISSN (P): 2790-7406, (E): 2790-7414
Volume 5, Issue 4 (Oct-Dec 2024)



Review Article



Emerging Paradigms in Exercise-Based Neuro-Physiotherapy for Holistic Motor and Cognitive Rehabilitation in Parkinson's Disease

Sumbal Imama¹ and Zulekha Zameer²

¹Institute of Physical Therapy, The University of Lahore, Lahore, Pakistan

²Kausar Abdulla Malik School of Life Sciences, Forman Christian College (A Chartered University), Lahore, Pakistan

ARTICLE INFO

Keywords:

Parkinson's Disease, Neurodegenerative Disorder, Neuro-Physiotherapy, Multimodal Exercise Programs

How to Cite:

Imama, S., & Zameer, Z. (2024). Emerging Paradigms in Exercise-Based Neuro-Physiotherapy for Holistic Motor and Cognitive Rehabilitation in Parkinson's Disease: Neuro-Physiotherapy in Parkinson's Disease. *THE THERAPIST (Journal of Therapies & Rehabilitation Sciences)*, 5(04), 02-10. <https://doi.org/10.54393/tt.v5i04.241>

*Corresponding Author:

Sumbal Imama
Institute of Physical Therapy, The University of
Lahore, Lahore, Pakistan
imamasumbal@gmail.com

Received date: 5th November, 2024

Acceptance date: 25th December., 2024

Published date: 31st December, 2024

ABSTRACT

Parkinson's Disease (PD) is a progressive neurodegenerative disorder that affects motor and non-motor functions, including cognitive, emotional and autonomic systems, severely impacting quality of life. The motor symptoms of PD are successfully treated by traditional physiotherapy, but such treatments often fail to address the complexity and variety of PD. Advancements in exercise-based neuro-physiotherapy are reviewed, with a focus on innovative and multimodal approaches combining motor and cognitive rehabilitation. Technology-driven interventions like virtual reality, robotics and AI add real-time feedback and personalized care to therapy, while cognitive strategies like dual-task training and mindfulness practice address cognitive impairments. Comprehensive benefits of multimodal exercise programs that include aerobic, strength and flexibility exercises are targeted to achieve both physical and mental health. Comparative analysis of traditional, emerging and multimodal approaches shows their strengths and weaknesses and highlights the need for tailored interventions. Future directions are directed at longitudinal research, a combination of pharmacological and surgical treatments, and the use of biomarkers and AI to design personalized therapy to enhance outcomes and quality of life of PD patients.

INTRODUCTION

Parkinson's Disease (PD) is a slowly progressive neurodegenerative disorder with motor symptoms including tremor, rigidity, bradykinesia and postural instability resulting from the loss of dopamine-producing neurons in the brain. In addition to these motor symptoms, PD also has a range of non-motor symptoms including cognitive impairment, depression, anxiety, sleep disorders and autonomic dysfunction [1]. Non-motor symptoms present so severely that they significantly impair patients' quality of life and complicate disease management, and therefore require holistic and adaptable therapeutic approaches [2, 3]. Innovative, technology-driven and multimodal approaches to address the complex nature of PD have been emerging trends in neuro-physiotherapy.

Balanced training and cognitive stimulation are being done with the use of virtual reality and gaming technologies, while dual-task training combines motor and cognitive exercises to improve overall functionality [4]. A more robust approach to rehabilitation is multimodal exercise programs, that include aerobic, strength, and flexibility exercises. Also, AI-powered tools are adding a layer of personalized therapy plans, making it possible for therapists to consider individual patient's needs for their treatment [5]. The purpose of this review is to explore new neuro-physiotherapy approaches in PD that may enhance motor and cognitive rehabilitation. This paper reviews the state of the science of PD rehabilitation, with a discussion of the limitations of traditional therapies and the benefits



of innovative solutions. It is anticipated that it will demonstrate how these advances can address the PD problem and improve patient outcomes.

Pathophysiology Basis of Parkinson's Disease

Parkinson's disease (PD) is a progressive neurodegenerative disorder of the motor system characterized by a progressive degeneration of dopamine-producing neurons in the substantia nigra. PD is a complex pathology that consists of several interrelated mechanisms leading to motor and non-motor symptoms [6].

Neurodegenerative Mechanism

The loss of dopaminergic neurons in the substantia nigra pars compacta of the basal ganglia is the hallmark feature of PD. Deficiency of dopamine is essential for smooth, coordinated movement and results in the characteristic motor symptoms of PD, including bradykinesia (slowness of movement), rigidity, resting tremor, and postural instability [7]. This loss of dopamine arises because the neurons that release dopamine into the striatum, the brain's main motor control center, undergo progressive degeneration [8].

Lewy Bodies and Protein Mis-folding

Another significant component of PD pathology is the appearance of additional Lewy bodies and abnormally formed protein lumps found in neurons. Primarily, they are made up of alpha-synuclein, a protein that functions in normal synaptic vesicle function and regular dopamine release. PD involves a mis-folding and aggregation of alpha-synuclein, which ultimately characterizes the formation of Lewy bodies [9]. Deposition of alpha-synuclein in other brain regions is thought to interfere with neuronal function and contribute to neuronal death or degeneration. Cognitive decline and psychiatric symptoms in PD patients result, over time, from the spread of these aggregates from the substantia nigra to other parts of the brain, including the cortex [10].

Mitochondrial Dysfunction and Oxidative Stress

Dysfunction of the mitochondria is critical to PD pathogenesis. The cells release energy for their host cell, and mitochondrial dysfunction implies reduced energy delivery to neurons, demanding high-energy neurons of the substantia nigra. Oxidative stress contributes by accumulating reactive oxygen species (ROS), which damage cellular components (lipids, proteins, and DNA). One reason neurons die or degenerate in PD cases is the inability of neurons to repair oxidative damage [11].

Neuro-Inflammation

Neuro-inflammation is another important pathological feature in PD. Exacerbation of neuronal damage is due to activated microglia (brain immune cells) and the release of pro-inflammatory cytokines. These neuro-inflammatory responses can be the cause as well as the consequence of neurodegeneration, constituting a vicious cycle of cellular damage. Neuro-inflammation has been associated with the progression of both motor and non-motor symptoms

[12, 13].

Genetic and Environmental Factors

Most PD cases are sporadic, but genetic mutations in specific genes, including LRRK2, PARK7, PINK1, and SNCA (encoding for alpha-synuclein), make one more prone to PD [14]. PD is also reported to be triggered by environmental factors like exposure to toxins like pesticides. The age of onset and severity of the disease may be influenced by genetic and environmental factors [1].

Neuroplasticity and Compensatory Mechanisms

One of the most important features in the progression of PD is the brain's attempt to compensate for dopaminergic neuron loss. Neuroplasticity is how the brain recruits other brain regions to take over lost functions [15]. However, these compensatory mechanisms stop working overtime as the disease progresses, and more severe symptoms become more apparent. Exercise enhances the formation of new neural connections and may slow down the progression of the disease [16].

Non-motor symptoms and Systemic Involvement

Although motor symptoms are the defining feature of PD, non-motor symptoms are often more associated with inducing disabling symptoms. Non-motor symptoms generally include cognitive decline, sleep disturbance, autonomic dysfunction (e.g., orthostatic hypotension), and psychiatric symptoms such as depression and anxiety. They are believed to involve widespread brain areas, including the cortex, the limbic system, and the brainstem [17]. Cognitive decline in PD is usually expressed as executive dysfunction (problems with planning and decision-making) and later dementia (problems with memory and reasoning) [18].

Motor Rehabilitation Through Exercise-Based Neuro-Physiotherapy

Management and improvement of movement require exercise-based motor rehabilitation. Body weight-supported treadmill training (BWSTT) is helpful for patients to train in walking, correct abnormal gait patterns, and increase walking speed and balance [19]. Doing these types of resistance exercises, such as with elastic bands or weights, will strengthen muscles and might slow the progression of PD when done regularly [20]. The methods increase motor skills like precision and strength, but they must be continued long-term to retain that benefit. Other therapies can be combined with these exercises to make them more effective. However, each patient's best intensity and duration still require more research [16, 21]. Using robotic aids and virtual reality (VR), people with PD are being helped with improving movement and balance using technology-assisted physiotherapy. VR creates interactive environments for balance and coordination training, while robotic devices, such as exoskeletons and treadmills, help with walking and good posture. However, these technologies are expensive or unsuitable for some patients, making therapy more engaging and effective. However, it is important to remember that traditional

exercises such as knee extensions, sit-to-stand, and backward walking remain important to improving strength, balance, and stability. Combined, these methods present a better approach to managing PD and a better quality of life for patients [16, 19]. It overviews key techniques such as body weight, body-supported treadmill training (BWSTT), resistance training, robotic assistance, virtual reality (VR) therapy, and dance therapy and describes their features, benefits, and limitations. Studies have been conducted

that were intended to enhance PD patients' strength, balance, gait, coordination, and flexibility, thereby contributing to perhaps achieving mobility and/or better quality of life. The table further addresses the challenges these therapies can encounter, for example, its cost, patient-specific, and others, and uses insights from recent research sources on these issues. A summary of various motor rehabilitation strategies for PD patients is shown (Table 10).

Table 1: Motor Rehabilitation Techniques in Parkinson's Disease

Technique	Key Features	Benefits	Limitations	Sources
Physical Therapy (PT)	Involves exercises to improve strength, flexibility, balance, and gait. Tailored to individual needs based on PD stage	Improves motor function, reduces rigidity, increases mobility, enhances balance and posture	May require frequent sessions for long-term improvement. Limited by physical space or resources	[20,21]
Gait Training	Focuses on walking improvements, including cadence, stride length, and reducing freezing of gait (FOG). Can use auditory or visual cues to aid movement	Reduces FOG episodes, improves walking speed and stability, decreases fall risk	May require continuous practice to maintain gains. Some patients may not respond to cues	[22,23]
Strength Training	Involves resistance exercises designed to build muscle mass and strength, often using weights or resistance bands	Enhances muscle strength, reduces rigidity, improves overall mobility, and reduces falls	Potential for overexertion or injury if not properly managed. Requires supervision	[24,25]
Dance and Rhythm Therapy	Dance styles (e.g., tango, ballet) or rhythmic exercises (e.g., drumming) that focus on coordination, posture, and rhythm	Improves coordination, balance, flexibility, and motor planning. Reduces motor symptoms and boosts mood	May be difficult for patients with advanced PD or severe rigidity. Accessibility issues	[26,27]
LSVT BIG Therapy	Amplitude-based therapy with exercises emphasizing large, exaggerated movements to overcome rigidity and bradykinesia	Increases motor amplitude, improves gait, reduces bradykinesia, enhances balance and overall mobility	Requires intensive and sustained effort over weeks/months. May be tiring for some patients	[28]
Cueing Techniques	Use of external auditory, visual, or tactile cues to help patients initiate and perform movements, especially during freezing episodes	Reduces freezing episodes, improves movement initiation and execution, enhances motor control	Not all patients respond equally to cues. May not be effective in severe PD cases	[29]
Aquatic Therapy	Exercises performed in a pool, using water resistance and buoyancy to reduce joint impact while enhancing strength, flexibility, and motor coordination	Enhances strength, balance, and flexibility. Provides a low-impact alternative to land-based exercises	Limited availability of facilities. May be challenging for those with severe mobility issues	[30]
Robotic-assisted Gait Training	Use of robotic devices or exoskeletons that assist in walking, helping patients relearn gait patterns through repetitive movements and support	Promotes neuroplasticity, improves gait, walking speed, reduces bradykinesia and fall risk	High cost, limited availability in clinics, may require training to use effectively	[31]
Treadmill Training	Walking exercises performed on a treadmill with or without body support. Often includes high-speed walking to improve gait and cardiovascular endurance	Improves cardiovascular fitness, walking speed, gait, and endurance. Reduces fall risk	May cause discomfort or fatigue, especially in advanced stages of PD. Requires careful monitoring	[32]

Neurofeedback	A technique that trains patients to self-regulate brain activity using real-time feedback on brainwave patterns. Often used to reduce tremors and improve focus	Reduces tremors, improves motor function, enhances focus and attention, potentially improves cognitive functions	Expensive, may require specialized equipment and trained clinicians. Limited long-term data	[33]
---------------	---	--	---	------

Cognitive Rehabilitation: Neuro-Physiotherapy's Dual Role

These cognitive problems memory loss, or trouble planning or multitasking in PD can make daily life harder and make movement worse [34]. Many cognitive challenges like focus, planning movements and remembering motor patterns can reduce the success of therapy [35]. This neuro-physiotherapy addresses the cognitive as well as the motor problems together. Often, it uses exercises that increase attention, memory, problem-solving, and movement tasks. Patients can work on both thinking and movement at the same time via techniques such as dual-tasking training or virtual reality. The combination of approaches improves independence and quality of life for people with PD [33, 36]. Advanced neuro-physiotherapy techniques for PD address both motor and non-motor symptoms (Table 2). Dual-task training involves movement and thinking tasks, and patients learn to multitask, pay attention, and become more independent [4, 37, 38]. Meditation and yoga are mindfulness practices that reduce anxiety and depression and increase emotional stability and movement [39-42]. In addition, these methods may also produce positive brain changes, such as increased grey matter in important areas. Although a promising but new method, neuro-feedback, which uses brain activity feedback to improve control has been used [43-46]. But these techniques look promising and further research is needed to validate their long-term benefits and develop effective treatment plans.

Table 2: Summary of cognitive rehabilitation strategies and their neurophysiological targets.

Technique	Key Features	Benefits	Limitations	Sources
Dual-task Training	Combines cognitive and motor tasks to improve overall function	Improves multitasking ability Enhances attention and executive function Promotes independence in daily activities	Optimal protocols still under research Challenging for advanced cognitive decline patients	[4, 37, 38]
Mindfulness Interventions	Incorporates mindfulness meditation and yoga to address psychological and motor symptoms	Reduces anxiety and depression Improves emotional stability and cognitive performance Enhances motor symptom management	Efficacy varies among individuals Long-term effects need more study	[39,41]
Neuro-feedback	Uses real-time brain activity feedback to improve self-regulation of neural processes	May improve motor control and reduce tremors Enhances cognitive and emotional regulation	Research in early stages Long-term effectiveness remains unclear	[43,44]

Role of Multimodal Exercise Programs

PD holistic rehabilitation combines exercises to assist in movement and general health. These programs combine aerobic exercises (heart disease, brain protection), strength training (muscle building and balance), and flexibility exercises (reduce stiffness and improve motion). Group-based or individualized programs are also available to patients. Social interaction and motivation are significant in group sessions, while an individual can focus on his needs and specific symptoms in the case of personalized plans. It is a case of some centers using both approaches. The best results come together. This combined method helps manage PD symptoms and improves quality of life [47]. Multimodal exercise programs have shown significant benefits for managing various conditions, including PD and cancer. In a randomized trial in Hong Kong, 138 PD patients were assigned to mindfulness yoga or stretching and resistance exercises. The yoga group had significant improvements in depression, anxiety,

motor function (MDS-UPDRS III scores reduced from 34.9 to 21.1), and quality of life [40]. For pancreatic cancer patients, a study of six older individuals undergoing chemotherapy found that five completed the program, showing increases or maintenance in lean mass (0.1%-4.4%), reduced fat mass (0.4%-8.6% in four patients), improved muscle strength (7.1%-75% in four to five patients), and better physical abilities like sit-to-stand and backward walking [48]. A trial in Spain with 44 cancer patients (mean age 63.46 years) also showed that those in a multimodal exercise program experienced lower fatigue, better functional capacity (SPPB scores), and reduced dependency (Barthel Index) over the control group [49]. These experiments reveal that both physical and mental health are enhanced by multimodal exercise programs for diverse patient populations.

Technology and Innovation in Neuro-Physiotherapy

The field of neuro-physiotherapy has begun to transform its approach to the rehabilitation of Parkinson's Disease

(PD) patients concerning technology. Technology offers personalized, engaging, adaptive therapies that improve motor and cognitive rehabilitation. Specific technologies applied directly include virtual reality (VR), rehabilitation robotics, wearable devices and sensors, artificial intelligence (AI) and machine learning, brain-computer interfaces (BCIs), and neuro-feedback, as well as integrating technology with traditional therapies[50-52].

Virtual Reality (VR)

VR is a valuable tool for PD patient's motor and cognitive rehabilitation. VR systems provide immersive environments where patients can practice movements, balance, and coordination tasks in a controlled and safe space. For example, on virtual trails or obstacle courses, walking will also improve gait and decrease the frequency of PD freezing episodes, a typical motor symptom of PD. Furthermore, VR systems include cognitive challenges, such as dual tasking (e.g., walking and solving puzzles), to train executive functioning and multi-tasking [4]. Studies have demonstrated that VR-based therapies can improve motor symptoms such as stride length, walking speed, and balance, as well as non-motor symptoms such as apathy and depression. VR also provides immediate feedback and engaging exercises, thus increasing patient motivation and adherence to therapy. In the advanced stages of PD, however, VR is likely to have difficulties in sensory integration and cognitive overload, so the effectiveness of VR at this stage of PD needs to be investigated[49].

Robotics in PD Rehabilitation

One of the significant roles played by robotic devices such as exoskeletons, robotic arms, and treadmills is to help PD patients regain functional independence. For example, robotic-assisted gait training (RAGT) assists patients in learning walking patterns again by simulating natural gait mechanics. These devices provide a form of high-intensity training that is also consistent, which helps improve postural control, decrease tremors, and reduce bradykinesia (slow movement) [52]. Additionally, robotic systems can enable genuine time assistance and resistance oriented to the patient's abilities to promote neuroplasticity and muscle strengthening. Biofeedback systems have also been integrated with robotic therapies, allowing PD patients to see their movements and chronic their progress. Although these devices are quite effective, their high cost and limited accessibility hinder their widespread use[53].

Wearable Devices and Sensors

Accelerometers, gyroscopes, and smartwatches have become wearable technologies that can help monitor and treat symptoms of PD. In real-time, these devices track movement and tremors as well as gait patterns so that the therapists can track the progression of symptoms and the effectiveness of their interventions. For instance, wearables can pick up on the freezing of gait episodes to deliver auditory or vibratory cues to patients to make it

through. Furthermore, wearable devices allow for ongoing monitoring in a nonclinical setting, providing important information about daily motor fluctuations and non-motor symptoms such as sleep disturbance and fatigue. Therefore, this data supports the customization of therapy plans and precision of care for PD patients[54].

Artificial Intelligence (AI) and Machine Learning

Analysis of patient data opens the doors of AI as a game changer in PD rehabilitation to personalize therapy plans. Machine learning algorithms can predict PD symptoms' progression, determine what works best for different individuals, and help select the most effective interventions. Therefore, AI-driven systems can analyze gait patterns based on the treadmill settings or determine the proper robotics assistance levels for specific therapy sessions. AI also helps make early diagnoses by analyzing imaging and biomarker data so early interventions can be started sooner. Additionally, it enables remote care via virtual coaching systems, which enable AI to provide real-time feedback on how exercise performance can be improved and thus recommend an adjustment, ensuring consistency and quality of therapy provided to PD patients in the home[52, 53].

Brain-Computer Interfaces (BCIs) and Neuro-feedback

The solutions to PD symptoms are promising solutions given by BCIs and neuro-feedback systems. PD patients can, with BCIs, use brain signals to control assistive devices like robotic arms, giving independence to people who are severely motor impaired. Instead, Neuro-feedback trains patients to control brain activity that could reduce tremors, mood, and cognitive functioning. Specifically, these systems are helpful for non-motor symptoms in PD, depression, and anxiety, which are very common in PD. In addition to modulating brain activity, neuro-feedback may also improve executive functions, including attention and memory, which are impaired in PD patients[43].

Combining Technology with Traditional Therapies

Although technological innovations may bring many benefits, their combination with mainstream physiotherapy practice provides the best results for PD patients. For example, with strength training exercises, VR and robotics can be combined, or wearables can provide biofeedback to a person during yoga or mindfulness sessions. The hybrid approach guarantees that motor and non-motor symptoms are treated holistically[55].

Comparative Analysis of Approaches

Each traditional, emerging, and multimodal approach has strengths and challenges (Table 3). It is simple, cost-effective, and easy to use but not flexible and scalable. In addition to VR and AI, other emerging approaches offer personalized treatments, real-time feedback, and greater engagement, but are expensive, complex, and have limited long-term efficacy data. Combining different methods in multimodal approaches can improve accuracy and engagement but is resource-intensive, more challenging

to implement, and may be overwhelming for some patients. The high costs of diagnosis and treatment, limited access, and patient compliance are challenges that call for a balanced approach that combines traditional and innovative methods and is appropriate for the patient and resources[56].

Table 3: Comparative Analysis of Neuro-Physiotherapy Paradigms Based On Key Parameters

Variables	Traditional Approaches	Emerging Approaches	Multimodal Approaches
Efficacy	Effective for specific tasks but lacks contextual adaptability	Highly effective with personalized treatments but limited long-term data	Comprehensive and robust but may overwhelm some patients
Cost	Low Cost	High cost due to advanced technology	Very high cost
Accessibility	Widely accessible in most settings	Limited accessibility in resource-constrained settings	Variable accessibility depending on resources and expertise
Engagement	Limited patient engagement	High patient engagement with real-time feedback	Enhanced engagement through diverse methods
Flexibility	Low flexibility	Moderate flexibility	Highly flexible for various needs
Resource Requirements	Minimal resource requirements	High resource requirements	Very high resource requirements

CONCLUSIONS

Parkinson's disease (PD) is a complex motor and non-motor challenge that requires holistic and creative rehabilitation. Motor symptoms respond well to traditional physiotherapy, but the need remains for more adaptive physiotherapy for cognitive impairments and the progressive nature of the disease. Among emerging technologies, virtual reality, robotics, artificial intelligence, and multimodal exercise programs promise personalized, engaging, and complete care. These advancements combine motor and cognitive rehabilitation to provide physical and mental health. Nevertheless, resource demands and accessibility remain intractable. Longitudinal research, integrating therapies with pharmacologic and surgical treatments, and using biomarkers and AI to personalize interventions are the future directions for this work. A balanced, patient-centred approach to PD can significantly improve outcomes and the quality of life of PD patients.

Authors Contribution

Conceptualization: SI

Methodology: SI, ZZ

Formal analysis: SI, ZZ

Writing, review and editing: SI, ZZ

All authors have read and agreed to the published version of the manuscript

Conflicts of Interest

The authors declare no conflict of interest.

Source of Funding

The authors received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Kumaresan M, Khan S. Spectrum of non-motor symptoms in Parkinson's disease. *Cureus*. 2021 Feb, 13(2).
- [2] Radad K, Moldzio R, Krewenka C, Kranner B, Rausch WD. Pathophysiology of non-motor signs in Parkinson's disease: some recent updating with brief presentation. *Exploration of Neuroprotective Therapy*. 2023 Feb, 3(1): 24-46.
- [3] Brunelli F, Torosantucci L, Gelmetti V, Franzone D, Grünwald A, Krüger R, Arena G, Valente EM. PINK1 protects against staurosporine-induced apoptosis by interacting with Beclin1 and impairing its pro-apoptotic cleavage. *Cells*. 2022 Feb, 11(4): 678.
- [4] Zheng Y, Meng Z, Zhi X, Liang Z. Dual-task training to improve cognitive impairment and walking function in Parkinson's disease patients: A brief review. *Sports Medicine and Health Science*. 2021 Dec, 3(4): 202-6.
- [5] Mylius V, Maes L, Negele K, Schmid C, Sylvester R, Brook CS, Brugger F, Perez-Lloret S, Bansi J, Aminian K, Paraschiv-Ionescu A. Dual-task treadmill training for the prevention of falls in Parkinson's disease: rationale and study design. *Frontiers in rehabilitation sciences*. 2022 Mar, 2: 774658.
- [6] Dong-Chen X, Yong C, Yang X, Chen-Yu S, Li-Hua P. Signaling pathways in Parkinson's disease: molecular mechanisms and therapeutic interventions. *Signal transduction and targeted therapy*. 2023 Feb, 8(1): 73.
- [7] Dauer W, Przedborski S. Parkinson's disease: mechanisms and models. *Neuron*. 2003 Sep, 39(6):88 9-909.
- [8] Song J, Kim J. Degeneration of dopaminergic neurons due to metabolic alterations and Parkinson's disease. *Frontiers in aging neuroscience*. 2016 Mar, 8: 65.
- [9] Calabresi P, Mechelli A, Natale G, Volpicelli-Daley L, Di Lazzaro G, Ghiglieri V. Alpha-synuclein in Parkinson's disease and other synucleinopathies: from overt neurodegeneration back to early synaptic dysfunction. *Cell death & disease*. 2023 Mar, 14(3): 176.
- [10] Rosen BI. Alpha-Synuclein Multimerization Is Dependent on Structural Characteristics of Repeated Ktkegv Regions. St. John's University (New

- York), 2021.
- [11] Zhou C, Wang L, Cheng W, Lv J, Guan X, Guo T *et al*. Two distinct trajectories of clinical and neurodegeneration events in Parkinson's disease. *NPJ Parkinson's Disease*. 2023 Jul, 9(1): 111.
 - [12] Çınar E, Tel BC, Şahin G. Neuroinflammation in parkinson's disease and its treatment opportunities. *Balkan medical journal*. 2022 Sep, 39(5): 318.
 - [13] Jurcau A, Andronie-Cioara FL, Nistor-Cseppento DC, Pascalau N, Rus M, Vasca E *et al*. The involvement of neuroinflammation in the onset and progression of Parkinson's disease. *International Journal of Molecular Sciences*. 2023 Sep, 24(19): 14582.
 - [14] Salles P, Tirapegui JM, Chaná-Cuevas P. Genetics of Parkinson's disease: Dominant forms and GBA. *Neurology Perspectives*. 2024 Jul, 4(3): 100153.
 - [15] Passaretti M, Cilia R, Rinaldo S, Rossi Sebastiano D, Orunesu E, Devigili G *et al*. Neurophysiological markers of motor compensatory mechanisms in early Parkinson's Disease. *Brain*. 2024 Nov, 147(11): 3714-26.
 - [16] Mak MK, Wong-Yu IS. Exercise for Parkinson's disease. *International review of neurobiology*. 2019 Jan, 147:1-44.
 - [17] Kallás MS, Nóbrega KH, dos Santos Figueiredo I, Padilha GB, de Rezende Pinto L. Oral dentures and quality of life related to Parkinson disease: An integrative review. *Research, Society and Development*. 2024 Jun, 13(6): e13713644657-.
 - [18] Andersson S, Josefsson M, Stiernman LJ, Rieckmann A. Cognitive decline in Parkinson's disease: a subgroup of extreme decliners revealed by a data-driven analysis of longitudinal progression. *Frontiers in Psychology*. 2021 Sep, 12: 729755.
 - [19] Takeuchi N, Izumi SI. Rehabilitation with poststroke motor recovery: a review with a focus on neural plasticity. *Stroke research and treatment*. 2013, 2013(1): 128641.
 - [20] Savoie FA, Benoit S, Riesco E, Tanguay A. Long-term impact of a community-based adapted boxing program on physical functioning and quality of life of individuals with Parkinson's disease. *NeuroRehabilitation*. 2024 Jan: 1-2.
 - [21] Burtcher J, Moraud EM, Malatesta D, Millet GP, Bally JF, Patoz A. Exercise and gait/movement analyses in treatment and diagnosis of Parkinson's Disease. *Ageing Research Reviews*. 2024 Jan, 93: 102147.
 - [22] Taniguchi S, D'cruz N, Nakagoshi M, Osaki T, Nieuwboer A. Determinants of impaired bed mobility in Parkinson's disease: Impact of hip muscle strength and motor symptoms. *NeuroRehabilitation*. 2022 Jan, 50(4): 445-52.
 - [23] Khobkhun F, Hollands M, Tretriluxana J, Srivanitchapoom P, Richards J, Ajjimaporn A. Benefits of task-specific movement program on en bloc turning in Parkinson's disease: A randomized controlled trial. *Physiotherapy Research International*. 2022 Oct, 27(4): e1963.
 - [24] Strand KL, Cherup NP, Totillo MC, Castillo DC, Gabor NJ, Signorile JF. Periodized resistance training with and without functional training improves functional capacity, balance, and strength in Parkinson's disease. *The Journal of Strength & Conditioning Research*. 2021 Jun, 35(6): 1611-9.
 - [25] Claus I, Muhle P, Czechowski J, Ahring S, Labeit B, Suntrup-Krueger S. Expiratory muscle strength training for therapy of pharyngeal dysphagia in Parkinson's disease. *Movement Disorders*. 2021 Aug, 36(8): 1815-24.
 - [26] Karpodini CC, Dinas PC, Angelopoulou E, Wyon MA, Haas AN, Bougiesi M *et al*. Rhythmic cueing, dance, resistance training, and Parkinson's disease: A systematic review and meta-analysis. *Frontiers in neurology*. 2022 Aug, 13: 875178.
 - [27] Moratelli JA, Alexandre KH, Boing L, Swarowsky A, Corrêa CL, de Guimarães AC. Dance rhythms improve motor symptoms in individuals with Parkinson's disease: a randomized clinical trial. *Journal of Dance Medicine & Science*. 2022 Mar, 26(1): 1-6.
 - [28] Eldemir S, Eldemir K, Saygili F, Ozkul C, Yilmaz R, Akbostancı MC *et al*. The effects of standard and modified LSVT BIG therapy protocols on balance and gait in Parkinson's disease: A randomized controlled trial. *Brain and Behavior*. 2024 Mar, 14(3): e3458.
 - [29] Magdi A, Sayed Ahmed AM, Elsayed E, Ahmad R, Ramakrishnan S, Gabor MG. Effects of cueing techniques on gait, gait-related mobility, and functional activities in patients with parkinson's disease: a systematic review and meta-analysis. *Physical Therapy Reviews*. 2021 Mar, 26(3): 188-201.
 - [30] Carroll LM, Morris ME, O'Connor WT, Clifford AM. Community aquatic therapy for Parkinson's disease: an international qualitative study. *Disability and Rehabilitation*. 2022 Jul, 44(16): 4379-88.
 - [31] Nedergård H, Arumugam A, Sandlund M, Bråndal A, Häger CK. Effect of robotic-assisted gait training on objective biomechanical measures of gait in persons post-stroke: a systematic review and meta-analysis. *Journal of Neuroengineering And Rehabilitation*. 2021 Dec, 18: 1-22.
 - [32] Tsai SY, Tai CH, Lee YY. Exploring Potential Predictors of Treadmill Training Effects in People With Parkinson Disease. *Archives of Physical Medicine and Rehabilitation*. 2024 Mar, 105(3): 525-30.

- [33] Anil K, Hall SD, Demain S, Freeman JA, Ganis G, Marsden J. A systematic review of neurofeedback for the management of motor symptoms in Parkinson's disease. *Brainsciences*. 2021 Sep, 11(10): 1292.
- [34] Rodríguez-Hernández M, Polonio-López B, Corregidor-Sánchez AI, Martín-Conty JL, Mohedano-Moriano A, Criado-Álvarez JJ. Can specific virtual reality combined with conventional rehabilitation improve poststroke hand motor function? A randomized clinical trial. *Journal of Neuro Engineering and Rehabilitation*. 2023 Apr, 20(1): 38.
- [35] Gilat M, Ginis P, Zoetewij D, De Vleeschhauwer J, Hulzinga F, D'Cruz N et al. A systematic review on exercise and training-based interventions for freezing of gait in Parkinson's disease. *npj Parkinson's Disease*. 2021 Sep, 7(1): 81.
- [36] Lee EY. Memory Deficits in Parkinson's Disease Are Associated with Impaired Attentional Filtering and Memory Consolidation Processes. *Journal of Clinical Medicine*. 2023 Jul, 12(14): 4594.
- [37] Oliveira de Carvalho A, Murillo-Rodríguez E, Rocha NB, Carta MG, Machado S. Physical exercise for Parkinson's disease: Clinical and experimental evidence. *Clinical Practice and Epidemiology in Mental Health*. 2018 Mar, 14(1).
- [38] Fang C, Lv L, Mao S, Dong H, Liu B. Cognition deficits in Parkinson's disease: mechanisms and treatment. *Parkinson's disease*. 2020: 2020(1); 2076942.
- [39] Hofheinz M, Mibs M, Elsner B. Dual task training for improving balance and gait in people with stroke. *The Cochrane Database of Systematic Reviews*. 2016: 2016(10).
- [40] Bender AD, Filmer HL, Naughtin CK, Dux PE. Dynamic, continuous multitasking training leads to task-specific improvements but does not transfer across action selection tasks. *NPJ Science of Learning*. 2017 Dec, 2(1): 14.
- [41] Bogosian A, Hurt CS, Vasconcelos e Sa D, Hindle JV, McCracken L, Cubi-Molla P. Distant delivery of a mindfulness-based intervention for people with Parkinson's disease: the study protocol of a randomised pilot trial. *Pilot and Feasibility Studies*. 2017 Dec, 3: 1-3.
- [42] van der Heide A, Meinders MJ, Speckens AE, Peerbolte TF, Bloem BR, Helmich RC. Stress and mindfulness in Parkinson's disease: clinical effects and potential underlying mechanisms. *Movement Disorders*. 2021 Jan, 36(1): 64-70.
- [43] Kwok JY, Choi EP, Wong JY, Lok KY, Ho MH, Fong DY et al. A randomized clinical trial of mindfulness meditation versus exercise in Parkinson's disease during social unrest. *NPJ Parkinson's disease*. 2023 Jan, 9(1): 7.
- [44] He S, Fang W, Wu J, Lv H, Zhang J, Wang T et al. Whether mindfulness-guided therapy can be a new direction for the rehabilitation of patients with Parkinson's disease: a network meta-analysis of non-pharmacological alternative motor-/sensory-based interventions. *Frontiers in psychology*. 2023 Sep, 14: 1162574.
- [45] Krylova M, Skouras S, Razi A, Nicholson AA, Karner A, Steyerl D et al. Progressive modulation of resting-state brain activity during neurofeedback of positive-social emotion regulation networks. *Scientific reports*. 2021 Dec, 11(1): 23363.
- [46] Yu L, Long Q, Tang Y, Yin S, Chen Z, Zhu C, et al. Improving emotion regulation through real-time neurofeedback training on the right dorsolateral prefrontal cortex: evidence from behavioral and brain network analyses. *Frontiers in Human Neuroscience*. 2021 Mar, 15: 620342.
- [47] Ventegodt S, Gringols M, Merrick J. Clinical holistic medicine: holistic rehabilitation. *The Scientific World Journal*. 2005, 5(1): 280-7.
- [48] Luo, H., Galvão, D.A., Newton, R.U., Tang, C.I., Spry, N. and Taaffe, D.R. Multimodal Exercise in Older Patients with Advanced Pancreatic Cancer Undergoing First-Line Chemotherapy: A Case Series Examining Feasibility and Preliminary Efficacy. *European Journal of Cancer Care*. 2023: 2023(1), 1179848.
- [49] Fernandez-Rodriguez EJ, Sanchez-Gomez C, Mendez-Sanchez R, Recio-Rodriguez JI, Puente-Gonzalez AS, Gonzalez-Sanchez J et al. Multimodal physical exercise and functional rehabilitation program in oncological patients with cancer-related fatigue A randomized clinical trial. *International Journal of Environmental Research and Public Health*. 2023 Mar, 20(6): 4938.
- [50] Huang Z, Meng L, Bi X, Xie Z, Liang W, Huang J. Efficacy and safety of robot-assisted deep brain stimulation for Parkinson's disease: a meta-analysis. *Frontiers in Aging Neuroscience*. 2024 May, 16: 1419152.
- [51] Parekh AD, Shaikh OA, Manan S, Al Hasibuzzaman M. Artificial intelligence (AI) in personalized medicine: AI-generated personalized therapy regimens based on genetic and medical history. *Annals of Medicine and Surgery*. 2023 Nov, 85(11): 5831-3.
- [52] Olawade DB, Wada OZ, Odetayo A, David-Olawade AC, Asaolu F, Eberhardt J. Enhancing mental health with Artificial Intelligence: Current trends and future prospects. *Journal of medicine, surgery, and public health*. 2024 Apr: 100099.
- [53] Sierra M SD, Garcia A DE, Otálora S, Arias-Castro MC,

Gómez-Rodas A, Munera M, Cifuentes CA. Assessment of a robotic walker in older adults with Parkinson's disease in daily living activities. *Frontiers in Neurorobotics*. 2021 Dec, 15: 742281.

- [54] Kehnemouyi YM, Coleman TP, Tass PA. Emerging wearable technologies for multisystem monitoring and treatment of Parkinson's disease: a narrative review. *Frontiers in Network Physiology*. 2024 Feb, 4: 1354211.
- [55] Das J, Barry G, Walker R, Vitorio R, Morris R, Stuart S. The integration of technology into a home-based visuo-cognitive training intervention for people with Parkinson's: Is the future digital?. *Plos one*. 2023 Jun, 18(6): e0285100.
- [56] Masinde R, Barasa D, Mandillah L. Effectiveness of using multimodal approaches in teaching and learning listening and speaking skills. 2023.